

Technical Report No. 19-03

Fish and Water Quality Monitoring at the Fort Knox Mine, 2018

by

Justin M. Burrows and Chad E. Bear



March 2019

Alaska Department of Fish and Game

Division of Habitat



Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the *Système International d'Unités* (SI), are used without definition in the following reports by the Divisions of Habitat, Sport Fish, and Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, Technical Reports and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

Weights and measures (metric)		General		Mathematics, statistics	
centimeter	cm	Alaska Administrative Code	AAC	<i>all standard mathematical signs, symbols and abbreviations</i>	
deciliter	dL	all commonly accepted abbreviations	e.g., Mr., Mrs., AM, PM, etc.	alternate hypothesis	H_A
gram	g	all commonly accepted professional titles	e.g., Dr., Ph.D., R.N., etc.	base of natural logarithm	e
hectare	ha	at	@	catch per unit effort	CPUE
kilogram	kg	compass directions:		coefficient of variation	CV
kilometer	km	east	E	common test statistics	(F, t, χ^2 , etc.)
liter	L	north	N	confidence interval	CI
meter	m	south	S	correlation coefficient (multiple)	R
milliliter	mL	west	W	correlation coefficient (simple)	r
millimeter	mm	copyright	©	covariance	cov
		corporate suffixes:		degree (angular)	$^\circ$
Weights and measures (English)		Company	Co.	degrees of freedom	df
cubic feet per second	ft ³ /s	Corporation	Corp.	expected value	E
foot	ft	Incorporated	Inc.	greater than	>
gallon	gal	Limited	Ltd.	greater than or equal to	\geq
inch	in	District of Columbia	D.C.	harvest per unit effort	HPUE
mile	mi	et alii (and others)	et al.	less than	<
nautical mile	nmi	et cetera (and so forth)	etc.	less than or equal to	\leq
ounce	oz	exempli gratia (for example)	e.g.	logarithm (natural)	ln
pound	lb	Federal Information Code	FIC	logarithm (base 10)	log
quart	qt	id est (that is)	i.e.	logarithm (specify base)	log ₂ , etc.
yard	yd	latitude or longitude	lat or long	minute (angular)	'
		monetary symbols (U.S.)	\$, ¢	not significant	NS
Time and temperature		months (tables and figures): first three letters	Jan, ..., Dec	null hypothesis	H_0
day	d	registered trademark	®	percent	%
degrees Celsius	°C	trademark	™	probability	P
degrees Fahrenheit	°F	United States (adjective)	U.S.	probability of a type I error (rejection of the null hypothesis when true)	α
degrees kelvin	K	United States of America (noun)	USA	probability of a type II error (acceptance of the null hypothesis when false)	β
hour	h	U.S.C.	United States Code	second (angular)	"
minute	min	U.S. state	use two-letter abbreviations (e.g., AK, WA)	standard deviation	SD
second	s			standard error	SE
Physics and chemistry				variance	
all atomic symbols				population sample	Var var
alternating current	AC				
ampere	A				
calorie	cal				
direct current	DC				
hertz	Hz				
horsepower	hp				
hydrogen ion activity (negative log of)	pH				
parts per million	ppm				
parts per thousand	ppt, ‰				
volts	V				
watts	W				

TECHNICAL REPORT NO. 19-03

**FISH AND WATER QUALITY MONITORING AT THE
FORT KNOX MINE, 2018**

By

Justin M. Burrows and Chad E. Bear
Division of Habitat, Fairbanks

Alaska Department of Fish and Game
Division of Habitat
1300 College Rd, Fairbanks, Alaska, 99701

March 2019

Cover: Fyke Net in the Developed Wetlands, May 11, 2018. Photograph by Chad E. Bear.

Technical Reports are available through the Alaska State Library, Alaska Resources Library and Information Services (ARLIS) and on the Internet: http://www.adfg.alaska.gov/index.cfm?adfg=habitat_publications.main. This publication has undergone editorial and peer review.

Note: Product names used in the publication are included for completeness but do not constitute product endorsement. The Alaska Department of Fish and Game does not endorse or recommend any specific company or their products.

*Alvin G. Ott, Justin M. Burrows, and Chad E. Bear
Alaska Department of Fish and Game, Division of Habitat
1300 College Rd., Fairbanks, AK 99701-1599, USA*

This document should be cited as:

Burrows, J.M. Bear, C.E. 2019. Fish and Water Quality Monitoring at the Fort Knox Mine, 2018. Alaska Department of Fish and Game, Technical Report No. 19-03, Fairbanks, Alaska.

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

If you believe you have been discriminated against in any program, activity, or facility please write:

ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK 99811-5526

U.S. Fish and Wildlife Service, 4401 N. Fairfax Drive, MS 2042, Arlington, VA 22203

Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW MS 5230, Washington DC 20240

The department's ADA Coordinator can be reached via phone at the following numbers:

(VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648,

(Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

For information on alternative formats and questions on this publication, please contact:

ADF&G Division of Sport Fish, Research and Technical Services, 333 Raspberry Road, Anchorage AK 99518 (907) 267-2375

Table of Contents

List of Tables	ii
List of Figures	ii
Acknowledgements.....	ii
Executive Summary	1
Introduction.....	1
Methods.....	5
Water Quality.....	5
Fish.....	6
Results and Discussion	7
Water Supply Reservoir, Water Quality	7
Stilling Basin, Arctic Grayling	14
Water Supply Reservoir, Arctic Grayling.....	14
Arctic Grayling Spawning (Timing, Temperature, and Fry Presence).....	15
Arctic Grayling Catches and Metrics.....	20
Water Supply Reservoir, Burbot.....	23
Literature Cited	27
References.....	28
Appendix 1. A Summary of Mine Development with Emphasis on Biological Factors..	30
Appendix 2. Water Quality Data, from the Fort Knox Water Supply Reservoir (WSR), April 2018.	36
Appendix 3. Population estimates of Arctic Grayling in the Fort Knox Water Supply Reservoir (WSR), 1995-2017.	39
Appendix 4. Arctic Grayling Growth in the Fort Knox Water Supply Reservoir (WSR), 2017-2018.	40
Appendix 5. Population Estimate of Burbot ($\geq 400\text{mm}$) in the Fort Knox Water Supply Reservoir (WSR), 2001-2017.	41

List of Tables

Table 1. Winter (October 1 to April 30) water use from the WSR, 1997 to 2018.....	3
---	---

List of Figures

Figure 1. Aerial photograph of the Fort Knox Gold Mine WSR, tailings facility and pit – water supply reservoir in lower part of photo and the tailings dam and impoundment in the upper Fish Creek valley, photograph provided by FGMI.....	1
Figure 2. Pond F outlet channel in 2000 (left photo) and in 2010 (right photo).....	2
Figure 3. Spillway in water supply reservoir in late April 2018.....	2
Figure 4. Map of Fort Knox Water Supply Reservoir water quality sample sites, April 5, 6 2018 and spring 2018 fyke net sites.....	5
Figure 5. WSR water temperature profiles, April 2018.....	7
Figure 6. WSR dissolved oxygen (mg/L) profiles, April 2018.....	8
Figure 7. WSR dissolved oxygen percent saturation (%) profiles, April 2018.	8
Figure 8. Winter water column dissolved oxygen in Polar Bay 2001-2018.....	9
Figure 9. Discharge point for non-contact water from dewatering wells around the Fort Knox open pit.....	9
Figure 10. Average, maximum, and minimum late winter water column DO concentrations at Site 2 in the WSR, 1998-2018 (excluding 2000, 2011, 2012, and 2014).	10
Figure 11. WSR pH profiles, April 2018.....	11
Figure 12. WSR specific conductance ($\mu\text{S}/\text{cm}$) profiles, April 2018.	12
Figure 13. WSR oxidation reduction potential (mV) profiles, April 2018.....	13
Figure 14. Pond F outlet channel in 2000 (left photo) and in 2010 (right photo).....	14
Figure 15. A hydraulic jump (about 1 m high) formed in the channel connecting Ponds D and E that was a limiting obstacle to the upstream movement of fish May 23, 2018.....	15
Figure 16. Arctic grayling upstream of 1m high hydraulic jump on May 23, 2018.	16
Figure 17. The catch per unit of effort of Arctic grayling (# of fish/hr) in the lower fyke net in the developed wetlands, 2018.	17
Figure 18. Peak daily water temperatures in Pond F outlet channel in spring 2018.....	18
Figure 19. Peak daily water temperatures in Pond F outlet channel in spring 2010 to 2018.	18
Figure 20. The percent of the Arctic grayling females handled that were categorized as: not ripe, ripe, or spent; 2018.	19
Figure 21. Estimates of the Arctic grayling population (fish ≥ 200 mm) in the WSR, with 95% Confidence Intervals, 1995-2017.....	20
Figure 22. Number of new fish ≥ 200 mm that entered the population but would have been too small to mark in the previous year (based on growth of marked fish), 2002-2018.	21
Figure 23. Length frequency distribution of Arctic grayling in Spring 1995, 2016, 2017, and 2018.....	22
Figure 16. Average growth of marked Arctic grayling before the WSR (1995) and in summers 2014, 2015, 2016, and 2017.....	22
Figure 25. Length frequency of burbot in the WSR and developed wetlands, 2017.....	23
Figure 26. Catch per unit of effort (fish/trap day) of burbot in the WSR, 1996-2007 and 2012-2018.....	23

Figure 27. Catch per unit of effort (fish/trap day) of large burbot (≥ 400 mm) in the WSR, 1996-2007 and 2012-2018. 24

Figure 28. Burbot (≥ 400 mm) population estimates \pm 95% CI in the WSR, 2001-2005 and 2012-2017..... 25

Figure 29. Growth (median, maximum, and minimum) of marked burbot (≥ 400 mm) in the WSR, 2000-2004 and 2012-2017. 25

Acknowledgements

We thank Fairbanks Gold Mining Inc. (FGMI) (Bartly Kleven, Mark Huffington, Jennifer Pyecha, and Dave Stewart) for their continued support of our work to monitor fish and wildlife resources in the water supply reservoir, tributaries, and developed wetlands.

We would also like to thank Parker Bradley, Chelsea Clawson, and Maria Wessel for their assistance with fieldwork; and Al Ott and Audra Brase for constructive reviews of this report.

Executive Summary

Water Quality

Dissolved oxygen (DO) concentrations were measured in mid-April 2018 and for the fourth consecutive year DO concentrations were some of the highest seen since sampling began in 1998. Higher DO concentrations appear to be directly related to the discharge of non-contact mine water to the water supply reservoir (WSR).

Arctic Grayling in the Water Supply Reservoir

Sampling for Arctic Grayling was conducted from May 3-14, 2018 as the fish moved from the WSR to the developed wetlands. Spawning began around May 11, when peak water temperatures approached 4.0°C, and on May 14, 10% of the females handled were spent. This is slightly later than in 2017 when 75% of females were spent on May 15. Ripe females continued to enter the wetland complex from the WSR at the close of the sampling event on May 14.

Substantial recruitment of Arctic grayling was observed in spring 2018. 498 fish between 200 and 245 mm FL were newly captured and tagged.

The spring 2017 population estimate for Arctic grayling ≥ 200 mm from fork length (FL) was 7,141 fish (95% CI 6,210 to 8,072 fish).

Burbot in the Water Supply Reservoir

Burbot sampling was conducted during the Arctic grayling sampling event from May 3-14, 2018 and again from June 6-10, 2018. During the two sampling events, 213 burbot were captured in the developed wetlands and WSR. These fish ranged in size from 110 to 849 mm TL. 72 of the captured burbot were ≥ 400 mm TL and 12 had been previously captured in the 2017 sampling events.

The spring/fall 2017 population estimate of burbot (≥ 400 mm TL) was 201 fish (95% CI 124 – 278). This is the highest population estimate of burbot >400 mm since sampling began at Fort Knox in 1996.

Introduction

Fairbanks Gold Mining Incorporated (FGMI) began construction of the Fort Knox hard-rock gold mine in March 1995. The mine is in the headwaters of the Fish Creek drainage about 25 km northeast of Fairbanks, Alaska (Figure 1). The project includes an open-pit mine, mill, tailings impoundment, water supply reservoir (WSR), and related facilities. Construction of the WSR dam and spillway was completed in July 1996. In 2007, permits were issued for the construction, operation, and closure of a valley fill heap leach facility located in Walter Creek upstream of the tailings pond. In 2018, ore continued to be processed through the mill and through the Walter Creek valley fill heap leach. Exploration drilling continued in the vicinity of the existing open pit.



Figure 1. Aerial photograph of the Fort Knox Gold Mine WSR, tailings facility, pit – water supply reservoir in lower part of photo, and the tailings dam and impoundment in the upper Fish Creek valley, photograph provided by FGMI.

Rehabilitation of the disturbed habitats has been concurrent with mining activities, to the extent practicable, and natural revegetation of some areas has been rapid (Figure 2).



Figure 2. Pond F outlet channel in 2000 (left photo) and in 2010 (right photo).

When full, the WSR contains about 3,363 acre-feet (1.1 billion gallons) of water. Water levels have remained mostly constant since 1998, except during the winter in certain years when large amounts were removed for mining processes. In late April 2018, there was relatively high surface flow over the spillway (Figure 3) at the outlet of the WSR from the higher than average snowfall in 2018. No water removal has occurred since the winter of 2014-15 (Table 1).



Figure 3. Spillway in water supply reservoir in late April 2018.

In spring 2015, FGMI initiated a discharge of non-contact water from dewatering wells around the open pit. The discharge was authorized by a permit issued by the Alaska Department of Environmental Conservation. The discharge began in mid-March and has been continuous with some periodic shutdowns. The discharge water appeared to increase the dissolved oxygen

concentrations in the WSR in late winter. From October 1, 2017 to April 30, 2018, the total discharge was 806 acre feet of water from Outfall 001.

Table 1. Winter (October 1 to April 30) water use from the WSR, 1997 to 2018.

Year (October 1 to April 30)	Acre-Feet of Water Removed	Percent of Water Removed
1997/1998	660	19.6
1998/1999	605	18.0
1999/2000	577	17.2
2000/2001	1,464	43.5
2001/2002	320	9.5
2002/2003	337	10.0
2003/2004	279	8.3
2004/2005	716	21.3
2005/2006	659	19.6
2006/2007	299	8.9
2007/2008	1,176	35.0
2008/2009	817	24.3
2009/2010	1,167	34.7
2010/2011	187	5.6
2011/2012	59	1.8
2012/2013	1,837	54.6
2013/2014	1,399	41.6
2014/2015	104	3.1
2015/2016	0	0
2016/2017	0	0
2017/2018	0	0

Populations of Arctic grayling (*Thymallus arcticus*) and burbot (*Lota lota*) exist in the WSR, and both Arctic grayling and burbot inhabit the stilling basin below the WSR. Arctic grayling spawning occurs predominantly in the wetland complex between the WSR and the tailings dam. Burbot spawning, as documented by radio telemetry, likely occurs in Solo Bay where Solo Creek enters the WSR (Figure 4). Arctic grayling recruit into the stilling basin by going over the WSR spillway when the water is high, however burbot do not appear to leave the WSR by this method as no tagged burbot from the WSR have been caught in the stilling basin.

Fish monitoring has been performed annually at the Fort Knox mine and related facilities since 1992 and water quality sampling since 1997. This report summarizes fish and water quality data collected during 2018 and discusses these findings in relation to previous work. A chronology of events from 2011 to 2018, with emphasis on biological factors, is presented in Appendix 1. The chronology for the previous years 1992 to 2010 can be found in ADF&G Technical Report No. 10-5, *Arctic grayling and burbot studies at the Fort Knox Mine, 2010* (Ott and Morris, 2010). The References section contains a comprehensive list of all ADF&G/ADNR technical biological reports for Fort Knox.

Methods

Water Quality

Water quality sampling was conducted on April 5 and 6, 2018, when the WSR was ice covered (Figure 4). Temperature ($^{\circ}\text{C}$), dissolved oxygen (DO) concentration (mg/L), DO percent saturation (barometrically corrected), pH, specific conductance ($\mu\text{S}/\text{cm}$), oxidation reduction potential (ORP), and depth (m) were measured with a Hydrolab[®] Minisonde[®]5 water quality multiprobe connected to a Surveyor[®] 4 digital display unit. The multiprobe sensors were calibrated to suggested specifications prior to use. The DO sensor was calibrated using a saturated air method. Conductivity, ORP, and pH sensors were calibrated with fresh standard solutions. Winter water quality measurements were made at 1 m depth intervals from the surface to the bottom.




Figure 4. Map of Fort Knox WSR water quality sample sites, April 5 and 6, 2018 and spring 2018 fyke net sites.

Fish

Fish sampling methods included visual observations, fyke nets, angling, and hoop traps. Two fyke nets were set in the developed wetlands (Figure 4). On May 3, 2018, a fyke net was set in the lower end of the Fish Creek channel. A second fyke net was set in Pond F outlet on the same day. Fyke nets were checked every day to every other day until both nets were pulled on May 14. Fish from both nets were processed for the entire sampling period.

Burbot were caught in the fyke nets and in hoop traps baited with herring in the WSR. Seventeen hoop traps were used to capture burbot in the late spring (June 6 to June 10, 2018).

Arctic grayling were measured to fork length (FL, nearest mm), inspected for tags and spawning condition, and released. Burbot were measured to total length (nearest mm), inspected for tags, and released. Un-tagged Arctic grayling ≥ 200 mm and burbot ≥ 300 mm were marked with a numbered Floy® T-bar internal anchor tag. Abundance of Arctic grayling and burbot was estimated using Chapman's modification of the Lincoln-Petersen two-sample mark-recapture model (Chapman 1951) and variance was estimated (Seber 1982).

Results and Discussion

Water Supply Reservoir, Water Quality

Water quality data were collected prior to breakup on April 6, 2018 (Appendix 2). Ice thickness on the WSR was slightly less than 1 m at each sampling location. There was an average of 15cm of overflow at each of the sites that had been refrozen on the surface for approximately 7.5cm then had slushy water for the next 7.5cm on top of the permanently frozen reservoir ice. This water did not influence the water quality sampling below the permanent ice layer. Water temperature ranged from 0.11 °C to 2.02 °C and steadily increased with depth (Figure 5).




Figure 5. WSR water temperature profiles, April 2018.

Fish Creek Bay and Polar Bay had the highest recorded dissolved oxygen (DO) levels at 10.98 mg/L and 11.01 mg/L (Figure 6) of the six sample sites. These maximums were higher than years prior to 2015 and are primarily a result of the non-contact groundwater discharge that began in March 2015 approximately 1.6 km upstream from the Pond F outlet. Fish Creek Bay had the highest average DO concentration for the fourth year in a row with 7.10 mg/L followed by Polar Bay at 6.11mg/L. Similar results were found in 2015, 2016, and 2017. Prior to 2015, the average DO at Fish Creek Bay was recorded at 0.195 mg/L. DO generally decreased with depth at all sites measured. Temperature specific DO saturation (%) followed the same pattern with higher values in the upper WSR near the inlet of Fish Creek and Solo Creek and lower values near the lower end of the WSR near the spillway and furthest from the fresh water sources (Figure 7).




Figure 6. WSR dissolved oxygen (mg/L) profiles, April 2018.




Figure 7. WSR dissolved oxygen percent saturation (%) profiles, April 2018.

The DO concentrations during late winter of 2015 through 2018, at Polar Bay were much higher than those previously recorded starting in 2001. (Figure 8). The increase in DO concentrations is

probably due to the discharge of non-contact water to the Fish Creek valley upstream of the WSR (Figure 9).




Figure 8. Winter water column dissolved oxygen in Polar Bay 2001-2018.



Figure 9. Discharge point for non-contact water from dewatering wells around the Fort Knox open pit.

Average winter water column DO at Site 2, in the WSR, was above the 16 year running average for the fourth year in a row and the third highest on record, behind 2016 (Figure 10). This is likely

a result of the continual discharge of non-contact ground water into the Fish Creek Valley about 1.6 km upstream from Pond F outlet.




Figure 10. Average, maximum, and minimum late winter water column DO concentrations at Site 2 in the WSR, 1998-2018 (excluding 2000, 2011, 2012, and 2014).

At all sites, pH values were relatively similar throughout the water column compared to prior years. At most sites, pH remained relatively stable with depth before slightly decreasing (Figure 11).




Figure 11. WSR pH profiles, April 2018.

Specific conductance was similar among all six sites throughout the reservoir (Figure 12). Values generally decreased with depth until 7 meters then began to plateau and increase. Fish Creek Bay did not follow this trend and may have been due to the much shallower water depth and influence of non-contact ground water into Fish Creek.




Figure 12. WSR specific conductance ($\mu\text{S/cm}$) profiles, April 2018.

Oxidation reduction potential (ORP) measures the ability of a lake or river system to break down waste products, such as contaminants and/or dead biological material. The more oxygen is present in the water, the higher the ORP value. ORP was highest in Site 2 and lowest in Site 1 and Solo Bay (Figure 13). This is a similar trend to previous years. Values were uniform throughout the water column except for Site 2 where ORP sharply decreased below 19 m as water became anoxic.




Figure 13. WSR oxidation reduction potential (mV) profiles, April 2018.

Stilling Basin, Arctic Grayling

The stilling basin located immediately downstream of the WSR spillway is fed by groundwater, seepage flow, and surface flow. A narrow notch in the spillway was designed to accommodate surface water discharge from the WSR during winter, without forming aufeis. Aufeis in the spillway has never been observed since it was constructed. In spring 2018, a relatively large volume of water was flowing over the spillway from the higher than average winter snowfall in 2018. Fish sampling in the stilling basin was not conducted during summer 2018.

Water Supply Reservoir, Arctic Grayling

Arctic grayling were found throughout the Fish Creek drainage prior to construction of the WSR. However, fish were concentrated in flooded mine cuts in Last Chance Creek. The population appeared stunted: fish larger than 220 mm were rare, average annual growth was 9 mm, and size at maturity was small averaging 148 mm for males and 165 mm for females (Al Ott, Unpublished Data). Successful spawning was limited to inlets and outlets of the flooded mine cuts and upper Last Chance Creek. Flooding of the WSR inundated the inlets and outlets of mine cuts, thus eliminating this spawning habitat. Since flooding of the WSR, aufeis in Last Chance Creek has been substantial. Since 1998, successful spawning by Arctic grayling in Last Chance Creek has only been observed in 2004 and 2005.

Very few fry were captured or observed (< 10 fish) from 1996 through 1998 in the WSR and Last Chance Creek. In spring 1999, FGMI constructed an outlet channel to connect the developed wetland complex with the WSR (Figure 14). The outlet channel was constructed to bypass a perched pipe and provide fish access to potential spawning and rearing habitat in the wetland complex.



Figure 14. Pond F outlet channel in 2000 (left photo) and in 2010 (right photo).

Arctic grayling have successfully spawned in the wetland complex every year since 1999. However, in 2002, 2006, and 2007, substantial aufeis and resultant cold water temperatures in the

wetland complex, in addition to beaver dams, limited the access and availability of spawning habitat. In recent years, aufeis buildup has been relatively minor and more effective beaver management has been implemented, including the annual removal of dams throughout the wetlands complex.

Arctic Grayling Spawning (Timing, Temperature, and Fry Presence)

In 2018, aufeis in the wetland complex was minimal. Beaver dams in Pond D outlet and in the lower wetland complex had been rebuilt and were breached by ADF&G staff on May 3 and May 23 to allow fish passage. Arctic grayling had access to the wetland complex (Ponds E and F), but access was reduced further upstream by a 1 m high natural barrier in the channel connecting Ponds D and E (Figure 15). Several Arctic graylings were observed upstream of the vertical obstruction between Pond E and Horseshoe ponds on May 23 (Figure 16).



Figure 15. A hydraulic jump (about 1 m high) formed in the channel connecting Ponds D and E that was a limiting obstacle to the upstream movement of fish, May 23, 2018.



Figure 16. Arctic grayling upstream of 1m high hydraulic jump on May 23, 2018.

Fish were caught immediately after the fyke nets were set. Most of the fish caught were Arctic grayling, however 79 burbot were also captured as they moved upriver in Fish Creek. The Arctic grayling CPUE was low during the first few days of sampling as water was rising rapidly and the fyke nets were knocked over for short periods of time (Figure 17). On May 10, the lower fyke net had multiple holes from an aquatic mammal, so CPUE could not be calculated.




Figure 17. The catch per unit of effort of Arctic grayling (# of fish/hr) in the lower fyke net in the developed wetlands, 2018.

The 2018 daily peak water temperature taken at the Pond F outlet did not rise above 4°C until May 13 (Figure 18), which was slightly later than average but not as late as 2013 (Figure 19). The warmest spring occurred in 2016, while 2013 was the coldest. The difference in spawning time between 2013 and 2016 was almost a full month (Figure 19).




Figure 18. Peak daily water temperatures in Pond F outlet channel in spring 2018.




Figure 19. Peak daily water temperatures in Pond F outlet channel in spring 2010 to 2018.

Arctic grayling spawning typically takes place when water temperature is near 4°C, and this can be seen in the number of ripe fish increasing throughout the sampling period (Figure 20). On May 4, the first day of fish capture, 93% of the female Arctic grayling were categorized as not ripe. The number of not ripe fish steadily decreased throughout the sampling period to 6% on May 12, then slightly increased to 14% as new Arctic grayling moved from the WSR into the Fish Creek wetlands complex. On May 4, 7% of the Arctic grayling in the wetlands complex were classified as ripe. The number of ripe fish steadily increased to 77% on May 13. No fish were classified as spent during the first four of the five days of the sampling period, then spent fish peaked at 11% on May 12 and 13. The 2018 spawning condition and timing were similar to 2015 and 2016 but differed from 2017 when 32% of Arctic grayling were classified as spent by May 14 (Ott and Bradley 2017).




Figure 20. The percent of Arctic grayling females handled that were categorized as not ripe, ripe, or spent; 2018.

Arctic Grayling Catches and Metrics

The spring 2017 population abundance estimate for Arctic grayling ≥ 200 mm was 7,141 fish with a 95% CI of 6,176 to 8,018 fish (Figure 21). The population has been declining but relatively stable for the past six years but showed a sharp increase in 2017, which also is reflected in the increased recruitment rates.

The 2017 population abundance estimate of Arctic grayling in the WSR was calculated using spring 2017 as the mark event and spring 2018 as the recapture event. During the spring of 2018 there were 1,240 grayling captured ≥ 230 mm, of those 162 were recaptures from the spring 2017 tagging event. For the 2017 population estimate, Arctic grayling population length frequency distributions from 2017 and 2018 were compared to eliminate those fish handled in 2018 that would have been too small (<200 mm) to mark in spring 2017. 169 fish met these criteria and were not included in the population estimate.




Figure 21. Estimates of the Arctic grayling population (fish ≥ 200 mm) in the WSR, with 95% Confidence Intervals, 1995-2017.

Recruitment (defined as those fish entering the population ≥ 200 mm but smaller than fish marked the previous year) is highly variable among the sampling years but was highest in 2017 and declined in 2018 (Figure 22). Substantial recruitment events were observed in the spring of 2004,

2010, 2014, and 2017. A substantial recruitment event was defined as >300 fish encountered during a recapture sampling event that were not available for tagging based on size during the mark sampling event (typically fish between 200 and 240 mm).




Figure 22. Number of new fish ≥ 200 mm that entered the population but would have been too small to mark in the previous year (based on growth of marked fish), 2002-2018.

The 2018 length frequency distribution of Arctic grayling caught in the wetlands complex is presented in Figure 23. Data for 1995, 2016 and 2017 are also included in this figure. The length distribution frequency has increased since 1995. The 1995 data set was obtained before construction of the freshwater dam and reflects the stunted condition of the population at that time. Note that the entire population length distribution shifted lower in 2017 and 2018 than in 2016 with the addition of many new fish (406 in 2017 and 498 in 2018) between 200 and 245 mm long.




Figure 23. Length frequency distribution of Arctic grayling in Spring 1995, 2016, 2017, and 2018.

Average growth of Arctic grayling prior to development of the WSR ranged from 3 to 17 mm per year (Figure 24 and Appendix 4). After the WSR was flooded in 1995, annual growth for fish increased substantially. Average growth in summer 2014 was highest.




Figure 16. Average growth of marked Arctic grayling before the WSR (1995) and in summers 2014, 2015, 2016, and 2017.

Water Supply Reservoir, Burbot

In spring 2018, burbot were caught in fyke nets (n = 79) during May and hoop traps (n = 134) during June in the WSR. 213 total burbot were caught during spring 2018 sampling. Burbot ranged in size from 110 to 849 mm with an average length of 359 mm (Figure 25).




Figure 25. Length frequency of burbot in the WSR and developed wetlands, 2017.

The catch per unit of effort for hoop traps (number of burbot per hoop trap/24 hrs) fished in the WSR remains low as compared with higher catches that occurred following the flooding of the reservoir (Figure 26). Catches of smaller burbot were highest in 1998 (7.2 fish/day), but decreased quickly and have remained low, averaging 1.1 burbot/trap/day in 2018.




Figure 26. Catch per unit of effort (fish/trap day) of burbot in the WSR, 1996-2007 and 2012-2018.

Catches of large burbot (≥ 400 mm) followed a similar pattern with catches increasing after the WSR was flooded, decreasing the next several years, but have been stable or increasing the last six years (Figure 27).




Figure 27. Catch per unit of effort (fish/trap day) of large burbot (≥ 400 mm) in the WSR, 1996-2007 and 2012-2018.

In spring of 2018, we recaptured 17 burbot ≥ 400 mm. The number of burbot seen in spring 2018 was 213, of which 12 were recaptured fish seen in 2017. The estimated population of large burbot for summer 2017 was 201 fish (95% CI 124-278 fish) (Figure 28, Appendix 5). This was the highest ≥ 400 mm population estimate since 2001 with 2012 having the next closest of 193 fish. The total 2017 estimate of burbot ≥ 300 mm was 736 fish (95% CI: 450 to 1022), this was also an increase from the 2016 population estimate of 505 burbot. It should be noted that the 2018 sampling was performed in the spring, and in previous years sampling was performed in the fall. Therefore, the increased CPUE may be due to possible seasonal differences in the catchability of burbot.




Figure 28. Burbot (≥ 400 mm) population estimates \pm 95% CI in the WSR, 2001-2005 and 2012-2017.

Growth rates of burbot from 2000 to 2004 and from 2012 to 2017 are shown in Figure 29. Growth is from tagged fish (≥ 400 mm) marked or seen in the previous year and recaptured in the following year during the same season. Growth is highly variable and while the population appears stable, more large burbot were caught from 2000 to 2002 than from 2003 to 2017, and only one burbot in this size class was captured during the spring in both 2017 and 2018.




Figure 29. Growth (median, maximum, and minimum) of marked burbot (≥ 400 mm) in the WSR, 2000-2004 and 2012-2017.

Conclusion

Populations of Arctic grayling and burbot have been established in the Fort Knox WSR. The post-mining goal for the Arctic grayling population was set at 800 to 1,600 fish ≥ 200 mm. Our spring 2017 estimated population for Arctic grayling ≥ 200 mm was 7,141 fish which is an increase from the estimated 2016 population of 4,396. A goal for the burbot population was not set prior to construction, but a small population of burbot ≥ 400 mm is present in the WSR.

We plan to continue to work cooperatively with FGMI to collect data on fish resources and water quality in the WSR and to implement rehabilitation projects designed to increase fish and aquatic habitat values and terrestrial habitats. Active management of beaver populations within the developed wetlands appears to remain a critical component to the productive capacity of the wetland complex for Arctic grayling.

Literature Cited

- Chapman, D.G. 1951. Some practices of the hypergeometric distribution with applications to zoological censuses. University of California Publications in Statistics. 1:131-160.
- Ott, A.G. and W.A. Morris. 2010. Arctic grayling and burbot studies at the Fort Knox mine, 2010. Alaska Department of Fish and Game Technical Report 10-05. Division of Habitat. Juneau.
- Seber, G.A.F. 1982. The estimation of animal abundance. Charles Griffin & Company LTD.
- Ott, A.G. 2019. Alaska Department of Fish and Game, Habitat Biologist, Personal Communication.
- Ott, A.G. and P.T. Bradley. 2017. Fish and water quality monitoring at the Fort Knox Mine, 2017. Alaska Department of Fish and Game Technical Report 17-10. Division of Habitat. Fairbanks.
- Ott, A.G., H.L. Scannell, and P.T. Bradley. 2016. Aquatic biomonitoring at Red Dog Mine, 2015. Alaska Pollution Discharge Elimination System Permit No. AK00038652 (Modification #1). Alaska Department of Fish and Game Technical Report 16-01. Division of Habitat. Fairbanks.

References

- Ott, A.G. and P.T. Bradley. 2016. Arctic grayling and burbot studies at the Fort Knox mine, 2016. Alaska Department of Fish and Game Technical Report 16-09. Division of Habitat. Fairbanks.
- Ott, A.G., P.T. Bradley, and H.L. Scannell. 2015. Arctic grayling and burbot studies at the Fort Knox mine, 2015. Alaska Department of Fish and Game Technical Report 15-07. Division of Habitat. Juneau.
- Ott, A.G., W.A. Morris, H.L. Scannell, and P.T. Bradley. 2014. Arctic grayling and burbot studies at the Fort Knox mine, 2014. Alaska Department of Fish and Game Technical Report 14-08. Division of Habitat. Juneau.
- Ott, A.G., W.A. Morris, H.L. Scannell, and P.T. Bradley. 2013. Arctic grayling and burbot studies at the Fort Knox mine, 2013. Alaska Department of Fish and Game Technical Report 13-05. Division of Habitat. Juneau.
- Ott, A.G., W.A. Morris, and H.L. Scannell. 2012. Arctic grayling and burbot studies at the Fort Knox mine, 2012. Alaska Department of Fish and Game Technical Report 12-09. Division of Habitat. Juneau.
- Ott, A.G. and W.A. Morris. 2011. Arctic grayling and burbot studies at the Fort Knox mine, 2011. Alaska Department of Fish and Game Technical Report 10-06. Division of Habitat. Juneau.
- Ott, A.G. and W.A. Morris. 2009a. Arctic grayling and burbot studies at the Fork Knox mine, 2009. Alaska Department of Fish and Game Technical Report 09-05. Division of Habitat. Juneau.
- Ott, A.G. and W.A. Morris. 2009b. Arctic grayling and burbot studies at the Fork Knox mine, 2008. Alaska Department of Fish and Game Technical Report 09-01. Division of Habitat. Juneau.
- Ott, A.G. and W.A. Morris. 2007. Arctic grayling and burbot studies at the Fork Knox mine, 2007. Alaska Department of Natural Resources Technical Report 07-01. Office of Habitat Management and Permitting. Juneau.
- Ott, A.G. and W.A. Morris. 2006. Arctic grayling and burbot studies at the Fork Knox mine, 2006. Alaska Department of Natural Resources Technical Report 06-05. Office of Habitat Management and Permitting. Juneau.
- Ott, A.G. and W.A. Morris. 2005b. Arctic grayling and burbot studies at the Fork Knox mine, 2005. Alaska Department of Natural Resources Technical Report 05-06. Office of Habitat Management and Permitting. Juneau.
- Ott, A.G. and W.A. Morris. 2005a. Arctic grayling and burbot studies at the Fork Knox mine, 2004. Alaska Department of Natural Resources Technical Report 05-01. Office of Habitat Management and Permitting. Juneau.
- Ott, A.G. and W.A. Morris. 2003. Arctic grayling and burbot studies at the Fork Knox mine, 2003. Alaska Department of Natural Resources Technical Report 03-09. Office of Habitat Management and Permitting. Juneau.

References (continued)

- Ott, A.G. and W.A. Morris. 2002b. Arctic grayling and burbot studies in the Fork Knox water supply reservoir, stilling basin, and developed wetlands, 2002. Alaska Department of Fish and Game Technical Report 02-06. Habitat and Restoration Division. Juneau.
- Ott, A.G. and W.A. Morris. 2002a. Arctic grayling and burbot studies in the Fork Knox water supply reservoir and developed wetlands, 2001. Alaska Department of Fish and Game Technical Report 02-1. Habitat and Restoration Division. Juneau.
- Ott, A.G. and W.A. Morris. 2001. Arctic grayling and burbot studies in the Fort Knox water supply reservoir and developed wetlands. Alaska Department of Fish and Game Technical Report 01-2. Habitat and Restoration Division. Juneau.
- Ott, A.G. and W.A. Morris. 2000. Fish use of the Fort Knox water supply reservoir and developed wetlands. Alaska Department of Fish and Game Technical Report 00-1. Habitat and Restoration Division. Juneau.
- Ott, A.G. and W.A. Morris. 1999. Fish use of the Fort Knox water supply reservoir 1995-1998. Alaska Department of Fish and Game Technical Report 99-2. Habitat and Restoration Division. Juneau.
- Ott, A.G. and P. Weber Scannell. 1998. Fisheries use and water quality in the Fort Knox mine water supply reservoir. Alaska Department of Fish and Game Technical Report 98-1. Habitat and Restoration Division. Juneau.
- Ott, A.G. and A.H. Townsend. 1997. Fisheries use of the Fort Knox water supply reservoir 1996. Alaska Department of Fish and Game Technical Report 97-2. Habitat and Restoration Division. Juneau.
- Ott, A.G. and P. Weber Scannell. 1996. Baseline fish and aquatic habitat data for Fort Knox mine 1992 to 1995. Alaska Department of Fish and Game Technical Report 96-5. Habitat and Restoration Division. Juneau.
- Ott, A.G., P. Weber Scannell, and A.H. Townsend. 1995. Aquatic habitat and fisheries studies upper Fish Creek, 1992-1995. Alaska Department of Fish and Game Technical Report 95-4. Habitat and Restoration Division. Juneau.
- Weber Scannell, P. and A.G. Ott. 1994. Aquatic habitat of Fish Creek before development of the Fort Knox gold mine 1992-1993. Alaska Department of Fish and Game Technical Report 94-5. Habitat and Restoration Division. Juneau.
- Weber Scannell, P. and A.G. Ott. 1993. Aquatic habitat study, upper Fish Creek drainage, with an emphasis on Arctic grayling (*Thymallus arcticus*): baseline studies 1992. Alaska Department of Fish and Game Technical Report 93-4. Habitat and Restoration Division. Juneau.

Technical Reports summarizing field work can be found on the Alaska Department of Fish and Game, Division of Habitat Web Page:

http://www.adfg.alaska.gov/index.cfm?adfg=habitat_publications.main

Appendix 1. A Summary of Mine Development with Emphasis on Biological Factors

2011

- February 9, ADF&G provided input to ADNR on the environmental audit to be conducted in summer 2011. We identified several possible fish and wildlife enhancement projects originally recommended by Buell and Moody (2005).
- March 4, the ACOE issued a permit (POA-1992-574-M19) authorizing construction of the modified dam raise and expansion of the Tailings Storage Facility (TSF).
- April and May, several Plan of Operations amendments were issued by ADNR for work associated with the TSF, waste rock dumps, powerline, topsoil storage, and dewatering.
- May 2, ADF&G provided input to ADNR on the reclamation and closure plan for Fort Knox. Emphasis was on maintaining the existing developed wetland complex downstream of the TSF.
- Our spring sample event for Arctic grayling and burbot ran from May 9 to 24. We caught 1,194 Arctic grayling and 117 burbot in a fyke net set in the WSR.
- The estimated spring 2010 Arctic grayling population was 4,346 fish > 200 mm long and was an increase from the 2009 estimate of 3,223. Recruitment of new fish in spring 2011 was strong with 198 new fish < 230 mm marked.
- Arctic grayling spawned in the wetland complex from Pond D downstream. Beavers had not rebuilt the dams in the wetland complex.
- A constructed osprey nesting platform adjacent to the main pump house in the WSR was occupied in spring – one chick was seen in August. An active raven nest was observed on the rock cut near the freshwater dam.
- Water began flowing over the spillway on May 27, water had not reached the spillway since winter 2009/2010.
- June 2, ADF&G provided written comments on the Ft. Knox and True North environmental audit proposals.
- July 19, FGMI pumped about 10,440 gallons of water from the “801 Pond” downstream – environmental staff were notified and pumping was immediately stopped – water from the “801 Pond” is supposed to be pumped back into sump below the TSF.
- August 4, ADNR informed us of planned changes at Fort Knox including expansion of the heap leach facility from 160 to 300 million tons, the need for a ADEC permit to discharge non-contact water, and the long-term need for a permit and water treatment plant for closure.

Appendix 1 (continued)

2011

- September 13, ADNR approved the drilling of two monitoring wells in the headwaters of Victoria Creek. The purpose of these monitoring wells is to ensure water in Victoria Creek is not impacted by the increased elevation of tailings in the Pearl Creek drainage.
- September 28, we met with FGMI to discuss plans to discharge non-contact water from the Fort Knox pit to the WSR.

2012

- Our spring sample event (Arctic grayling and burbot) began on May 7 and ended on May 30. The estimated spring 2011 Arctic grayling population was 7,378 fish \geq 200 mm long which was an increase of 3,032 from the 2010 estimate. Recruitment of new fish in spring 2012 was strong with 111 new fish $<$ 230 mm marked.
- We caught 140 burbot (175 to 950 mm long) in spring 2012 in hoop traps and fyke nets.
- Arctic grayling spawned throughout the wetland complex, including the upper portion of Channel C, in spring 2012. Beavers had not rebuilt the dams in the wetland complex.
- A constructed osprey nesting platform adjacent to the main pump house in the WSR was occupied in spring 2012.
- Water was flowing over the spillway when we began sampling in the spring of 2012 – water was still overflowing in late October.
- July 13, ADF&G provided input to ADEC on the APDES draft permit for discharge of non-contact water. The discharge point has been changed to the old Fish Creek channel just downstream of Ponds A and B. The ADEC permit was issued on August 15, 2012.
- September 27, ADF&G confirmed that a culvert in the road down the Fish Creek valley had been removed. In our trip report to FGMI, we recommended some additional civil work to ensure that the discharge water stays on the north side of the valley.

2013

- February 20, FGMI received a Notice of Violation from the ACOE for the unauthorized discharge of fill material into 0.28 acres of wetlands.
- March 1, ADF&G informed FGMI that their 2012 Annual Report was extremely well done and FGMI's report was distributed to all habitat offices in the state.
- March 11, the ACOE issued an After-the-Fact authorization covering the 0.28 acres of wetland fill.

Appendix 1 (continued)

2013

- April 25, water quality data (temperature, dissolved oxygen, etc.) were collected in the WSR under ice cover.
- May 4, the ADNR transmitted comments on the December 2012 reclamation and closure plan.
- Our spring sample event (Arctic grayling and burbot) began on May 20 and ended on June 10. The estimated spring 2012 Arctic grayling population was 7,404 fish ≥ 200 mm long. Recruitment of new fish in spring 2013 was strong with 114 new fish < 230 mm marked.
- We caught 96 burbot (89 to 697 mm long) in spring 2013 in hoop traps and fyke nets.
- Arctic grayling spawned throughout the wetland complex, including the upper portion of Channel C, in spring 2013. Beavers had rebuilt the dams in the wetland complex, but the dams were notched to allow fish passage.
- A constructed osprey nesting platform adjacent to the main pump house in the WSR was occupied in spring 2013.
- Water was not flowing over the spillway when we began sampling, but by May 27 water had begun to flow out of the WSR and over the spillway.
- June 25, we observed Arctic grayling fry (numerous) in the upper portion of Channel C. Very few fry were observed in Pond F and the Pond F outlet.
- October 14, ADF&G submitted comments on the Fort Knox 2013 reclamation plan – eight recommendations were made.
- November 27, ADF&G distributed the Fork Knox technical report for work done in 2013.

2014

- In early April, emails were exchanged to determine when Fish Creek was removed from the list of impaired waterbodies – it was on the 1992 list but was removed from the 1994 list because FGMI had bought out all the existing placer operations and was planning on building the freshwater dam.
- April, 2014, the decision was made not to collect winter water quality due to unsafe ice conditions and overflow.
- In spring 2014, we fished a fyke net in the developed wetlands just upstream of the WSR from April 29 until May 9 and then again from May 12 to 15. Arctic grayling spawned throughout the wetland complex in spring 2014. The only beaver dam present was in the upper end of C Channel.
- Our estimated population of Arctic grayling (> 200 mm) for spring 2013 was 6,675 – a slight reduction from the 2011 and 2012 estimates.
- Our estimated population of large burbot (≥ 400 mm) for spring 2013 was 80 – a substantial reduction from the spring 2012 estimate of 193.

- September 29, FGMI notified state agencies that the new Environmental Manager was Bartly Kleven.

Appendix 1 (continued)

2014

- September 4, we were notified that the road across Solo Creek had failed – FGMI will determine a proper fix – this is the second time the road has failed at the culvert crossing.
- September 26, the developed wetlands and lower Last Chance Creek were inspected. No beaver dams were observed in Ponds D and F and in lower Last Chance Creek (dams had been removed by FGMI during summer).
- October FGMI and ADF&G discussed a draft design for the Solo Creek culvert replacement, conducted a field inspection, and continued discussions to decide what remedial work will be done.
- October 28, ADF&G distributed the Fork Knox technical report for work done in 2014.
- November 12, FGMI submitted a permit application to replace the Solo Creek culvert. ADF&G had several questions regarding the culvert design specifications and FGMI addressed these questions and a permit was issued on November 20, 2014 to install the new 10-foot diameter pipe.

2015

- March 2, we conducted a field visit to observe the discharge point for non-contact mine water to the old Fish Creek channel which is dry, except for breakup and periods of heavy rain.
- FGMI initiated the discharge of non-contact water (about 250 gallons per minute) in mid-March and the discharge has been continuous except for a few shutdowns. The discharge was authorized by a permit issued by the ADEC.
- April 8 and 9, we collected water quality data in the WSR which was ice covered, high DO concentrations were found in Fish Creek Bay.
- April 17, we collected water quality data in the old Fish Creek channel downstream from where the non-contact mine water was being discharged and found very high DOs in the water – leading us to conclude that the discharge of non-contact mine water resulted in increased DOs in the WSR.
- Early May, we field inspected the culvert replacement in Solo Creek and concluded that it had been installed in accordance with the Fish Habitat Permit.
- Spring 2015, we fished a fyke net in the developed wetlands just upstream of the WSR from May 4 to 8 and then again from May 10 to 13. Arctic grayling spawned throughout the wetland complex in spring 2015.
- Our estimated population of Arctic grayling (≥ 200 mm) for spring 2014 was 5,841 – a slight reduction from the 2011 and 2012 estimates.

- Our estimated population of large burbot (≥ 400 mm) for spring 2014 was 175 – a substantial increase from the spring 2013, but with a large 95% CI.
- June 19 and July 23, we collected Arctic grayling fry in the wetland complex, average size on June 19 was 29.7 mm and on July 23 it was 57.3 mm.

Appendix 1 (continued)

2015

- June 19, we inspected the Last Chance culvert in the Gil Causeway. Material at the east end of the pipe has slumped and the road was blocked with cones and flagging.
- October 28, ADF&G distributed the Fork Knox technical report for work done in 2015.

2016

- March 29 and 31, water quality data were collected at six sites in the WSR, five of which have been sampled nearly annually since 1998. Average winter water column dissolved oxygen at Site 2 (middle of the WSR) was the highest on record and likely the result of the near continuous discharge of non-contact water into the old Fish Creek channel just upstream of the wetland complex.
- In spring 2016, we fished two fyke nets in the developed wetlands just upstream of the WSR and in Pond F from April 25 to May 4. Based on the fyke net catches, most of Arctic grayling spawned in the wetland complex downstream of Pond F.
- Our estimated population of Arctic grayling (≥ 200 mm) for spring 2015 was 5,947 – a slight increase from the 2014 estimate.
- Our estimated population of large burbot (≥ 400 mm) for spring 2015 was 92 - a substantial decrease from spring 2014.
- In early October, hoop traps fished in the WSR captured 26 burbot ranging in size from 200 to 630 mm long.
- October 12, we met with ADEC and FGMI to discuss plans to design and install a new water treatment plant just downstream of the tailings dam with an estimated discharge of 2,000 to 6,000 gallons per minute.
- October 28, we were notified by FGMI that beaver dams at Pond D outlet and downstream of Pond F had been removed.
- December 21, ADF&G sent a summary of our meeting on the new water treatment plant to FGMI.

2017

- April 12 and 19, water quality data were collected at six sites in the WSR, five of which have been sampled nearly annually since 1998. Average winter water column dissolved oxygen at Site 2 (middle of the WSR) was above the 15 year running average and the second highest on record, behind 2016.

- In spring 2017, we fished two fyke nets in the developed wetlands just upstream of the WSR and in Pond F from early May to May 18. Based on the fyke net catches and observations, most Arctic grayling spawned in the wetland complex downstream of Pond F.
- About 100 Arctic grayling adults were moved from the Pond F fyke net and released into Pond D upstream of a barrier. These fish successfully spawned in Pond D as fry were captured on June 29.
- Our estimated population of Arctic grayling (≥ 200 mm) for spring 2016 was 4,396, a decrease of about 1,500 fish from 2015.

Appendix 1 (concluded)

2017

- May 26, ADEC issued Waste Management Permit 2014DB002 (Modification #1).
- May 26, ADNR issued a permit amendment for the construction of the Barnes Creek heap leach.
- July 19, ADNR issued a Certificate of Approval to construct a dam for the Barnes Creek heap leach (#AK00315).
- October 12, a site visit was conducted to check on the status of beaver dams in the wetland complex that had been removed recently by FGMI.
- October 24, historic information was provided to FGMI on the status of Fish Creek and why it was taken off the impaired waterbody list in 1994.
- December 12, FGMI, ADF&G, ADNR, and ADEC met to discuss alternatives for tailings disposal, closure configuration for the tailing dam at elevation 1557, and a new water treatment plant.
- December 13, FGMI acquired a new parcel of land that contains an estimated 2.1 million ounces of gold.

2018

- March 14, ADNR approved a POA amendment request to replace the power line trail.
- April 3, 5, and 6, water quality data were collected at six sites in the WSR, five of which have been sampled nearly annually since 1998.
- May 3-May 14, two fyke nets were fished in the developed wetlands just upstream of the WSR and in Pond F.
- Our estimated population of Arctic Grayling (≥ 200 mm) for Spring 2017 was 7,141, which is an increase of 2,745 over 2016.
- Our estimated population of large burbot (≥ 400 mm) was 201 fish, which is an increase of 82 fish over 2016.
- October 9, 2018, the Pond D beaver dam was removed to allow the downstream movement of grayling into the WSR.

Appendix 2. Water Quality Data, from the Fort Knox Water Supply Reservoir (WSR), April 2018.

Site Number (Name)	Date	Depth (m)	Temperature (C)	% Saturation Dissolved Oxygen	Dissolved Oxygen (mg/L)	Conductivity (μ S/cm)	pH	ORP
1 (Middle WSR)	4/6/2018	1	0.6	45.8	6.27	153.2	7.26	395
		2	0.68	40.7	5.57	144.5	7.31	390
		3	1.43	37.1	4.97	142.3	7.17	394
		4	1.92	38.6	5.1	142.6	7.14	394
		5	2.3	38.2	5.01	141.5	7.08	397
		6	2.54	36.9	4.79	141	7.05	397
		7	2.76	35.3	4.57	141	7.1	395
		8	2.9	34.2	4.41	140.7	7.06	397
		9	2.99	33.2	4.26	140.8	7.04	397
		10	3.03	32	4.12	141.3	7.04	397
		11	3.11	30.3	3.88	141.7	7.06	395
		12	3.19	29.4	3.76	141.3	7.04	395
		13	3.2	28.8	3.68	141.4	7.07	395
		14	3.22	28.1	3.6	141.8	7.06	394
		15	3.24	26.8	3.43	141.9	7.06	395
		16	3.23	19.2	2.54	143.7	7.04	396
2 (WSR Near Dam)	4/6/2018	1	0.54	39.6	5.43	145.2	6.98	424
		2	0.71	39.4	5.4	142.9	7.01	423
		3	1.4	38.7	5.19	141.5	7.02	423
		4	1.9	38	5.03	141.3	7.02	423
		5	2.25	36.7	4.81	140.9	7.01	424
		6	2.57	35.7	4.65	139.9	7.04	423
		7	2.75	35.1	4.54	140	7.06	422
		8	2.85	35.3	4.56	140.1	7.04	423
		9	3.08	35.7	4.57	139.3	7.11	420
		10	3.13	34.8	4.45	139.4	7.06	423
		11	3.2	33.6	4.3	139.7	7.08	422
		12	3.27	32.4	4.13	139.8	7.07	422
		13	3.32	30.2	3.85	140	7.06	422
		14	3.36	25.6	3.26	140	7.04	422
		15	3.37	20	2.55	141.6	7	424
		16	3.39	15.5	1.98	143.4	6.99	423
		17	3.42	10.6	1.36	144	6.97	424

	18	3.45	7.6	0.97	146.4	6.96	424
	19	3.48	1.9	0.25	151.3	6.95	425
	20	3.48	0.3	0.04	157	6.95	425
	21	3.44	0.1	0.02	163.3	6.98	391
3 (Solo Bay)	4/6/2018						
	1	0.28	62	8.59	154.1	6.79	403
	2	0.47	59.4	8.21	151	6.92	399
	3	1.01	54.1	7.39	146.4	6.94	400
	4	1.8	45.4	6.08	143.7	6.95	401
	5	2.15	40.1	5.29	143.4	6.91	404
	6	2.29	38.6	5.05	143.6	6.93	404
	7	2.58	33.4	4.34	143.1	6.93	405
	8	2.61	17.2	2.3	145.7	6.93	404
	9	2.6	12.9	1.69	146.5	6.88	406
	10	2.6	7.8	1.02	147.5	6.9	405
	11	2.5	7.2	0.95	148.3	6.93	404
7 (Lower Last Chance Bay)	4/6/2018						
	1	0.24	46.1	6.38	146.7	6.81	417
	2	0.56	32.6	4.47	142.2	6.87	419
	3	1.15	35.4	4.78	142.8	6.92	419
	4	1.8	37.9	5.03	142.4	6.97	419
	5	2.13	38.2	5.03	142.8	6.99	419
	6	2.48	36	4.69	142.6	7.02	418
	7	2.55	23.5	3.06	143.8	6.95	420
11 (Polar Bay)	4/6/2018						
	1	0.16	79.3	11.01	164.9	7.07	411
	2	0.51	65	8.98	153.3	6.98	413
	3	1.1	50.4	6.83	150.1	6.97	413
	4	1.67	48.2	6.43	148.2	6.98	413
	5	1.98	47.6	6.29	148.4	6.98	413
	6	2.29	47.8	6.27	147.1	7.02	411
	7	2.46	47.3	6.17	147.6	7.02	413
	8	2.66	47.8	6.19	148.2	7.06	411
	9	2.74	47.6	6.16	148	7.08	410
	10	2.74	47.9	6.2	149.5	7.08	410
	11	2.84	45	5.83	152.5	7.05	410
	12	3.22	20.4	2.61	154.8	6.89	414
	13	3.51	3.7	0.47	165.6	6.84	416
12 (Fish Creek Bay)	4/6/2018						
	1	0.22	78.6	10.98	145.5	6.9	402
	2	0.45	70.4	9.73	149	6.99	400

	3	1.33	48.5	6.53	147	6.96	404
	4	1.56	36.6	4.89	150.8	6.9	406
	5	1.86	25	3.37	150.7	6.89	405

Appendix 3. Population estimates of Arctic Grayling in the Fort Knox Water Supply Reservoir (WSR), 1995-2017.

Year	Population Estimate ¹	95% Confidence Interval
1995 ²	4,358	
1996 ³	4,748	3,824-5,672
1996 ⁴	3,475	2,552-4,398
1998 ⁵	5,800	4,705-6,895
1999	4,123	3,698-4,548
2000	5,326	4,400-6,253
2001	5,623	5,030-6,217
2002	6,503	6,001-7,005
2003	6,495	5,760-7,231
2004	6,614	5,808-7,420
2005	7,926	6,759-9,094
2006	5,930	5,382-6,478
2007	4,027	3,620-4,433
2008	3,545	3,191-3,900
2009	3,223	2,896-3,550
2010	4,346	3,870-4,823
2011	7,378	6,616-8,141
2012	7,404	6,775-8,033
2013	6,675	6,217-7,333
2014	5,841	5,235-6,446
2015	5,947	5,111-6,783
2016	4,396	3,913-4,880
2017	7,141	6,176-8,018

¹Population estimates from 1995-1996 include fish ≥ 150 mm, in all other years fish ≥ 200 mm.

²In 1995, we used estimates from the ponds and creeks for the Arctic grayling population; a confidence interval was not applicable to the data set.

³The 1996 estimate was made with a capture and recapture event in summer 1996 using fyke nets.

⁴In 1996, Arctic grayling were captured with a boat-mounted electroshocker for both the capture and recapture events in fall 1996 by Sport Fish Division.

⁵Starting in 1998 through 2017 the population estimates were made using a mark event in the spring of the year of the estimate, and the recapture event in spring of the following year.

Appendix 4. Arctic Grayling Growth in the Fort Knox Water Supply Reservoir (WSR), 2017-2018.

Upper Limit (mm)	Average (mm)	Maximum (mm)	Minimum (mm)	Sample Size
210	32	38	27	3
220	18	29	5	9
230	23	40	9	18
240	18	33	3	33
250	15	35	0	20
260	13	27	0	16
270	9	25	0	16
280	9	22	0	13
290	3	9	0	10
300	5	15	0	12
310	3	8	0	7
320	4	10	0	3
330	0	0	0	1
340	0	0	0	0
350	0	0	0	0

Appendix 5. Population Estimate of Burbot ($\geq 400\text{mm}$) in the Fort Knox Water Supply Reservoir (WSR), 2001-2017.

Year	Population Estimate	95% Confidence Interval
2001	134	58-210
2002	131	63-199
2003	102	57-147
2004	86	44-128
2005	143	96-191
2006-2011 – no population estimates performed		
2012	193	95-290
2013	80	44-117
2014	175	44-305
2015	92	45-138
2016	119	65-173
2017	201	124-278